

16. The perimeter of the window is

$$2 = 2h + 2r + \pi r.$$

Rearranging this gives $h = 1 - r - \frac{1}{2}\pi r$.

The area of the rectangle is $2rh$. If we say that 1 unit area of glass lets through 1 unit of light, then the rectangle admits $2rh$ units of light. The area of the semicircle is $\frac{1}{2}\pi r^2$. Because tinted glass admits only half as much light as clear glass, the semicircle lets $\frac{1}{2}(\frac{1}{2}\pi r^2) = \frac{1}{4}\pi r^2$ units of light through. Hence, the window admits

$$L(r) = 2rh + \frac{1}{4}\pi r^2 = 2r \left(1 - r - \frac{1}{2}\pi r \right) + \frac{1}{4}\pi r^2 = 2r - 2r^2 - \pi r^2 + \frac{1}{4}\pi r^2 = 2r - 2r^2 - \frac{3}{4}\pi r^2$$


units of light. We wish to maximise $L(r)$.

Since $0 = L'(r) = 2 - 4r - \frac{3}{2}\pi r$, we must have $2 = (4 + \frac{3}{2}\pi)r$ and $r = \frac{2}{4 + \frac{3}{2}\pi} = \frac{4}{8 + 3\pi}$ metres. Thus

$$h = 1 - r - \frac{1}{2}\pi r = 1 - \left(1 + \frac{1}{2}\pi \right) r = 1 - \left(\frac{2 + \pi}{2} \right) \left(\frac{4}{8 + 3\pi} \right) = 1 - \frac{4 + 2\pi}{8 + 3\pi} = \frac{4 + \pi}{8 + 3\pi}$$

metres.

17. (a) f has a critical point at $x = 1$.
(b) f is increasing on $(0, 1)$. f is decreasing on $(1, \infty)$.
(c) Since $f''(x)$ exists and is non-zero for all $x \in (0, \infty)$, f is concave down everywhere.
(d)



18. (a) $G(x) = \frac{x^4}{4} + \frac{1}{2x^2}$ is an antiderivative of $g(x) = x^3 - \frac{1}{x^3}$, because

$$G'(x) = \frac{d}{dx} \left(\frac{x^4}{4} + \frac{1}{2}x^{-2} \right) = \frac{4x^3}{4} + \frac{1}{2}(-2x^{-3}) = x^3 - \frac{1}{x^3} = g(x).$$

- (b) $H(x) = \frac{4}{3} \sec 3x$ is an antiderivative of $h(x) = 4 \sec 3x \tan 3x$, because

$$H'(x) = \frac{d}{dx} \left(\frac{4}{3} \sec 3x \right) = \frac{4}{3} \left(\frac{d}{du} \sec u \right) \left(\frac{d}{dx} 3x \right) = \frac{4}{3} (\sec u \tan u) (3) = 4 \sec 3x \tan 3x = h(x)$$

by the chain rule.

- (c) $L(x) = \frac{1}{2}(e^x - e^{-x})$ is an antiderivative of $l(x) = \frac{1}{2}(e^x + e^{-x})$, because

$$L'(x) = \frac{d}{dx} \left(\frac{1}{2}(e^x - e^{-x}) \right) = \frac{1}{2}(e^x - (-1)e^{-x}) = l(x)$$

by the chain rule.

19. This is wrong because

$$\begin{aligned} \frac{d}{dx} (xe^x + 3 \cot 3x) &= (x)'e^x + x(e^x)' + 3 \left(\frac{d}{du} \cot u \right) \left(\frac{d}{dx} 3x \right) \\ &= e^x + xe^x + 3(-\operatorname{cosec}^2 u)(3) \\ &= e^x + xe^x - 9 \operatorname{cosec}^2 3x \end{aligned}$$

by the product rule and the chain rule.

20. (a) $\int \left(8t^3 - \frac{t^2}{2} + t \right) dt = 2t^4 - \frac{t^3}{6} + \frac{t^2}{2} + C.$

(b) $\int (\sec^2 \pi \theta) d\theta = \frac{1}{\pi} \tan \pi \theta + C.$

(c) $\int \left(x + \frac{1}{x} \right)^2 dx = \int \left(x^2 + 2 + \frac{1}{x^2} \right) dx = \frac{1}{3}x^3 + 2x - \frac{1}{x} + C.$